

Goat Anti-Human Factor IX (F.IX) Whole IgG from Antiserum (10 mg)

Catalogue: GAFIX-IG

Lot No: SAMPLE

Expiry date: **/****

Store at -10 to -20°C.

For Research Use Only ~ Not for use in diagnostic procedures.

Description of Factor IX

Factor IX (F.IX, Christmas Factor) is a vitamin K-dependant glycoprotein produced in the liver. Plasma concentration of F.IX is normally around 5 ug/ml (87 nM) in plasma. The biological importance of F.IX is demonstrated in Haemophilia B (Christmas disease), an X-linked congenital bleeding disease resulting from a quantitative (low activity and low antigen) or qualitative (low activity and normal antigen) defect in F.IX function.

In its proenzyme or zymogen form F.IX is a single chain molecule of 55,000 daltons. It contains two EGF-like domains and an amino-terminal domain containing 12 γ-carboxy-glutamic acid (Gla) residues. These Gla residues allow F.IX to bind divalent metal ions and participate in calcium-dependant binding interactions. The activation of F.IX occurs by limited proteolysis in the presence of calcium by activated factor XI (FXI^a) and/or by a complex of VII^a/tissue factor/phospholipid and activated Factor X between residues Arg¹⁴⁶-Ala¹⁴⁷ and between Arg¹⁸⁰-Val¹⁸¹. The terminal activated product in either case is F.IX^a_β, a two-chain enzyme consisting of a heavy chain (28,000 daltons), a light chain (18,000 daltons) and an activation peptide product of 11,000 daltons. F.IX can also be cleaved into inactive products by thrombin and by elastase.

The activity of F.IX^a_β in plasma is inhibited by antithrombin and this inhibition is accelerated 1000-fold in the presence of optimal concentrations of heparin¹⁻³.

References and Reviews

1. Lawson, JH, Mann KG; Cooperative Activation of Human F.IX by the Human Extrinsic Pathway of Coagulation; JBC 266 pp11317-11327, 10991.
2. Enfield DL, Thompson AR; Cleavage and Activation of Factor IX by Serine Proteases; Blood 64, pp 821-831, 1984.
3. Limentani SA, Furie BC, Furie B in **Hemostasis and Thrombosis**, 3rd Edition, eds. RW Colman, J Hirsh, VJ Marder and EW Salzman, pp. 94-108, J.B. Lippincott Co., Philadelphia PA,

USA, 1994.

Product Specifications

Description:

Vial containing 1.000 ml of whole IgG representing approximately 1 mL of antiserum. Total protein is 10 mg.

Format:

Whole IgG, clear liquid.

Host Animal:

Goat

Immunogen:

Human Factor IX purified from plasma.

Concentration:

IgG concentration is 10 mg/ml, determined by absorbance using an extinction coefficient ($E_{280}^{1\%}$) of 13.4.

Buffer:

10 mM HEPES, 0.15 M NaCl, 50% (v/v) glycerol, pH 7.4.

Storage:

Store between -10 and -20 °C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers. Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use.

Specificity:

This antibody is specific for F.IX as demonstrated by immunoelectrophoresis and ELISA.

Applications:

Suitable as a source of enriched antibodies to F.IX.

Neutralizing Activity:

68 Bethesda Units/ml IgG against normal plasma. One Bethesda unit/ml is defined as the amount of inhibitor that resulted in 50% residual F.IX activity after 2 hours at 37°C (Kasper CK *et al*, Thromb Diath Haemorrh **34**:869 1975).

Related Products

GAFIX-AP	Goat anti-Human FIX, affinity-purified IgG
GAFIX-HRP	Goat anti-Human FIX, IgG-peroxidase
GAFIX-APHRP	Goat anti-Human FIX, IgG-peroxidase
SAFIX-IG	Sheep anti-Human FIX, IgG from antiserum
SAFIX-AP	Sheep anti-Human FIX, affinity-purified IgG
SAFIX-APHRP	Sheep anti-Human FIX, APIgG-peroxidase
FIX-EIA	Paired antibody set for ELISA of FIX, 5 x 96 wells
FIX-DP	Human plasma deficient in FIX, immune depleted

Visit our site (www.enzymeresearch.co.uk) for further details.

Enzyme Research Laboratories Ltd
Glamorgan House . Charter Court . Enterprise Park
Swansea . SA7 9DD . UK.
Tel: +44 (0)1792 686070
Email: info@enzymeresearch.co.uk
Website: www.enzymeresearch.co.uk